

## MAGNETISM AND MATTER


### SECTION – A

Questions 1 to 10 carry 1 mark each.

1. The figure below shows the North and South poles of a permanent magnet in which  $n$  turn coil of area of cross – section  $A$  is resting, such that for a current  $I$  passed through the coil, the plane of the coil makes an angle  $\theta$  with respect to the direction of magnetic field  $B$ . If the plane of the magnetic field  $B$ .


The torque on the coil will be.

- (a)  $\tau = nIAB \cos\theta$   
 (b)  $\tau = nIAB \sin\theta$   
 (c)  $\tau = nIAB$




- (d) None of the above, since the magnetic field is radial

2. A uniform magnetic field exists in space in the plane of paper and is initially directed from left to right. When a bar of soft iron is placed in the field parallel to it, the lines of force passing through it will be represented by



3. A magnetic dipole moment of a bar magnet is a vector quantity directed:
- (a) upward at perpendicular bisector to the line joining to north pole and south pole  
 (b) from North pole to South pole  
 (c) downward at perpendicular bisector to the line joining to north pole and south pole  
 (d) from South pole to North pole
4. A toroid of  $n$  turns, mean radius  $R$  and cross-sectional radius  $a$  carries current  $I$ . It is placed on a horizontal table taken as  $x$ - $y$  plane. Its magnetic moment  $m$
- (a) is non-zero and points in the  $z$ -direction by symmetry.  
 (b) points along the axis of the toroid ( $m = m\phi$ ).  
 (c) is zero, otherwise there would be a field falling at  $\frac{1}{r^3}$  large distances outside the toroid.  
 (d) is pointing radially outwards.
5. The variation of magnetic susceptibility with the temperature of a ferromagnetic material can be plotted as

# CBSE ACADEMY PLUS



6. A long solenoid has 1000 turns per metre and carries a current of 1 A. It has a soft iron core of  $\mu_r = 1000$ . The core is heated beyond the Curie temperature,  $T_c$ .
- (a) The H field in the solenoid is (nearly) unchanged but the B field decreases drastically.  
(b) The H and B fields in the solenoid are nearly unchanged.  
(c) The magnetisation in the core reverses direction.  
(d) The magnetisation in the core does not diminishes.
7. A magnet of dipole moment  $M$  is aligned in equilibrium position in a magnetic field of intensity  $B$ . The work done to rotate it through an angle  $\theta$  with the magnetic field is  
(a)  $MB \sin \theta$                       (b)  $MB \cos \theta$                       (c)  $MB (1 - \cos \theta)$                       (d)  $MB (1 - \sin \theta)$
8. A magnetic needle suspended parallel to a magnetic field requires  $\sqrt{3}$  J of work to turn it through  $60^\circ$ . The torque needed to maintain the needle in this position will be  
(a)  $2\sqrt{3}$  J                      (b) 3 J                      (c)  $\sqrt{3}$  J                      (d)  $\frac{3}{2}$  J

In the following questions 9 and 10, a statement of assertion (A) is followed by a statement of reason (R). Mark the correct choice as:

- (a) Both assertion (A) and reason (R) are true and reason (R) is the correct explanation of assertion (A).  
(b) Both assertion (A) and reason (R) are true but reason (R) is not the correct explanation of assertion (A).  
(c) Assertion (A) is true but reason (R) is false.  
(d) Assertion (A) is false but reason (R) is true.


9. **Assertion (A):** The poles of magnet cannot be separated by breaking into two pieces.  
**Reason (R):** The magnetic moment will be reduced to half when a magnet is broken into two equal pieces.
10. **Assertion (A):** The ferromagnetic substances do not obey Curie's law.  
**Reason (R):** At Curie point a ferromagnetic substance start behaving as a paramagnetic substance.

## SECTION – B

Questions 11 to 14 carry 2 marks each.


11. Explain the following:  
(i) Why do magnetic lines of force form continuous closed loops?  
(ii) Why are the field lines repelled (expelled) when a diamagnetic material is placed in an external uniform magnetic field?
12. Define magnetic susceptibility of a material. Name two elements, one having positive susceptibility and the other having negative susceptibility. What does negative susceptibility signify?
13. The following figure shows the variation of intensity of magnetisation versus the applied magnetic field intensity,  $H$ , for two magnetic materials  $A$  and  $B$ :

# CBSE ACADEMY PLUS



- (a) Identify the materials *A* and *B*.  
 (b) Why does the material *B*, have a larger susceptibility than *A*, for a given field at constant temperature?


14. The diagrams given in the figure (a) and (b) show magnetic field lines (thick lines in the figure) wrongly. Point out what is wrong with them. Some of them may describe electrostatic field lines correctly. Point out which ones.



## SECTION – C

Questions 15 to 17 carry 3 marks each.

15. (i) A uniform magnetic field gets modified as shown below when two specimens *X* and *Y* are placed in it. Identify whether specimens *X* and *Y* are diamagnetic, paramagnetic or ferromagnetic.




- (ii) How is the magnetic permeability of specimen *X* different from that of specimen *Y*?

OR

Define the term magnetic permeability of a magnetic material. Write any two characteristics of a magnetic substance if it is to be used to make a permanent magnet. Give an example of such a material.

16. Two identical magnetic dipoles each of magnetic dipole moment  $2 \text{ Am}^2$  are placed with their axes perpendicular to each other with a distance of separation being  $r = 2 \text{ m}$  between them.




# CBSE ACADEMY PLUS

Find the magnetic field at the point that is midway along the distance of separation between the two dipoles.


OR

Depict the field-line pattern due to a current carrying solenoid of finite length.

- In what way do these lines differ from those due to an electric dipole?
- Why can't two magnetic field lines intersect each other?



Field lines of a current carrying solenoid



Field lines of an electric dipole

17. A short bar magnet of magnetic moment  $0.9 \text{ J/T}$  is placed with its axis at  $30^\circ$  to a uniform magnetic field. It experiences a torque of  $0.063 \text{ J}$ .
- Calculate the magnitude of the magnetic field.
  - In which orientation will the bar magnet be in stable equilibrium in the magnetic field?

## SECTION – D

Questions 18 carry 5 marks.


18. Derive an expression for magnetic field intensity due to a magnetic dipole at a point on its axial line.

## SECTION – E (Case Study Based Questions)

Questions 19 to 20 carry 4 marks each.

### 19. Gauss's Law for Magnetism

By analogy to Gauss's law of electrostatics, we can write Gauss's law of magnetism as  $\oint \vec{B} \cdot d\vec{s} = \mu_0 m_{\text{inside}}$  where  $\oint \vec{B} \cdot d\vec{s}$  is the magnetic flux and  $m_{\text{inside}}$  is the net pole strength inside the closed surface. We do not have an isolated magnetic pole in nature. At least none has been found to exist till date. The smallest unit of the source of magnetic field is a magnetic dipole where the net magnetic pole is zero. Hence, the net magnetic pole enclosed by any closed surface is always zero. Correspondingly, the flux of the magnetic field through any closed surface is zero.



- Consider the two idealised systems
  - a parallel plate capacitor with large plates and small separation and
  - a long solenoid of length  $L \gg R$ , radius of cross-section.

# CBSE ACADEMY PLUS

In (I)  $\vec{E}$  is ideally treated as a constant between plates and zero outside. In (ii) magnetic field is constant inside the solenoid and zero outside. These idealised assumptions, however, contradict fundamental laws as below

(a) case (I) contradicts Gauss's law for electrostatic fields.

(b) case (II) contradicts Gauss's law for magnetic fields.

(c) case (I) agrees with  $\oint \vec{E} \cdot d\vec{l} = 0$ .

(d) case (II) contradicts  $\oint \vec{H} \cdot d\vec{l} = I_{en}$ .

(ii) The net magnetic flux through any closed surface, kept in a magnetic field is

(a) zero (b)  $\mu_0/4\pi$  (c)  $4\pi\mu_0$  (d)  $4\mu_0/\pi$

(iii) A closed surface S encloses a magnetic dipole of magnetic moment  $2ml$ . The magnetic flux emerging from the surface is

(a)  $\mu_0 m$  (b) zero (c)  $2\mu_0 m$  (d)  $2m/\mu_0$

(iv) Which of the following is not a consequence of Gauss's law?

(a) The magnetic poles always exist as unlike pairs of equal strength.

(b) If several magnetic lines of force enter in a closed surface, then an equal number of lines of force must leave that surface.

(c) There are abundant sources or sinks of the magnetic field inside a closed surface.

(d) Isolated magnetic poles do not exist.

OR


(v) The surface integral of a magnetic field over a surface

(a) is proportional to mass enclosed (b) is proportional to charge enclosed

(c) is zero (d) equal to its magnetic flux through that surface.

## 20. Elements of the Earth's Magnetic Field

The earth's magnetic field at a point on its surface is usually characterised by three quantities: (a) declination (b) inclination or dip and (c) horizontal component of the field. These are known as the elements of the earth's magnetic field. At a place, angle between geographic meridian and magnetic meridian is defined as magnetic declination, whereas angle made by the earth's magnetic field with the horizontal in magnetic meridian is known as magnetic dip.



(i) In a certain place, the horizontal component of magnetic field is  $1/\sqrt{3}$  times the vertical component. The angle of dip at this place is

(a) zero (b)  $\pi/3$  (c)  $\pi/2$  (d)  $\pi/6$

(ii) The angle between the true geographic north and the north shown by a compass needle is called as

(a) inclination (b) magnetic declination (c) angle of meridian (d) magnetic pole.

(iii) The angles of dip at the poles and the equator respectively are

(a)  $30^\circ, 60^\circ$  (b)  $0^\circ, 90^\circ$  (c)  $45^\circ, 90^\circ$  (d)  $90^\circ, 0^\circ$

(iv) A compass needle which is allowed to move in a horizontal plane is taken to a geomagnetic pole. It

# CBSE ACADEMY PLUS

- (a) will become rigid showing no movement
- (b) will stay in any position
- (c) will stay in north-south direction only
- (d) will stay in east-west direction only.

**OR**

- (v) Select the correct statement from the following.
- (a) The magnetic dip is zero at the centre of the earth.
- (b) Magnetic dip decreases as we move away from the equator towards the magnetic pole.
- (c) Magnetic dip increases as we move away from the equator towards the magnetic pole.
- (d) Magnetic dip does not vary from place to place.

